



Management of Perinatal Care
Clinical Practice Guideline
MedStar Health

“These guidelines are provided to assist physicians and other clinicians in making decisions regarding the care of their patients. They are not a substitute for individual judgment brought to each clinical situation by the patient’s primary care provider-in collaboration with the patient. As with all clinical reference resources, they reflect the best understanding of the science of medicine at the time of publication but should be used with the clear understanding that continued research may result in new knowledge and recommendations”.

General Principles: The MedStar Health Obstetrical Service line endorses the ACOG Perinatal Care Guidelines and ACOG Committee Opinions as a source of guidance for clinical care of women throughout the MedStar Health system.

The following is a summary of clinical actions excerpted from:

- 1) American Academy of Pediatrics, American College of Obstetricians and Gynecologists. Guidelines for Perinatal Care. 8th ed. Elk Grove Village (IL): AAP; Washington, DC: American College of Obstetricians and Gynecologists; 2017. 693 pages. ISBN 9781934984967 (ACOG); ISBN 9781610020879 (AAP)
2) ACOG Committee Opinion Number 736: Optimizing Postpartum Care, May 2018
3) Immunization recommendations are from CDC guidelines, 2018; https://www.cdc.gov/vaccines/schedules/index.html
4) ACOG Committee Opinion 691. The citation is: Carrier screening for genetic conditions. Committee Opinion No. 691.American College of Obstetricians and Gynecologists. Obstet Gynecol 2017;129:e41–55.

Preconception care is acknowledged as an important component of high-quality prenatal care but is beyond the scope of this summary document. Additional information on this topic can be found in the source document

Prenatal Care:

Quality Goals:

- Improve the timeliness of prenatal care.
• Prenatal care within the first trimester or within 42 days of enrollment.
• Provide education and recommended screening and intervention.
• Monitor progression of pregnancy.
• Assess the well-being of the woman and her fetus.
• Early detection and intervention of high-risk factors.
• Complete 80% of expected prenatal visits. (ACOG recommends 14 visits).
• Decrease the incidence of smoking during pregnancy.
• Improve the frequency of appropriate testing during pregnancy.

Office visits

Frequency:

- Advise office visit at 8-10 weeks of pregnancy (or earlier if the patient is at risk for ectopic pregnancy).
• Every 4 weeks for first 28 weeks.
• Every 2 – 3 weeks until 36 weeks gestation.
• Every week after 36 weeks gestation.

Table with 3 columns: Initial Approval Date and Reviews, Most Recent Revision and Approval Date, Next Scheduled Review Date. Content includes dates from 2014 to 2022 and condition: Perinatal Care.

Frequency of visits is determined by individual needs and assessed risk factors.

First Prenatal Visit (8-10 weeks of pregnancy if first contact earlier)

Assessment at this visit should include:

- Initial history and physical.
- Family medical history.
- Genetic history.
- General exam to confirm pregnancy.
- Complete needs assessment.
- Preterm labor risk, education and prevention.
- Assess for tobacco, alcohol, drug use.
- Domestic violence screening.
- Screen for depression (current or historical) using a standardized screening tool.
- Obtain travel history with specific attention to areas where Zika virus is reported.
- Prescriptions: prenatal vitamins and iron supplementation as necessary.

Education and counseling

- Scope of care provided in the office and anticipated schedule of visits.
- Expected course of pregnancy.
- Counseling regarding specific complications.
- Discuss routine lab studies/testing.
- Discuss genetic counseling and available prenatal diagnostic testing (invasive and non-invasive) including non-invasive prenatal screening (NIPS) for aneuploidy testing, nuchal translucency (NT) sonogram and blood testing for First Trimester Screening (FTS).
- Schedule testing at appropriate gestational age
 - Nuchal translucency and FTS done at 11-13 weeks gestation
 - Cell free DNA, non-invasive prenatal screening (NIPS) can be done any time after 10 weeks and should be accompanied by appropriate counseling.
 - NIPS and invasive testing (chorionic villus sampling, amniocentesis) would typically apply to all women over 35 or otherwise at increased risk for aneuploidy, preceded by genetic counseling.
- Discuss high risk conditions.
- Education regarding: Labor and delivery, nutrition, exercise, working, air travel, routine dental care, tobacco use and smoke exposure, alcohol/drug consumption, over-the-counter medications, pets, etc.
- Practices to promote health maintenance such as use of safety restraints including lap and shoulder belts.
- Assess barriers to care (transportation, childcare issues, work schedule).
- Encourage maternity program enrollment and prenatal classes.
- Make needed referrals for medical conditions, substance abuse and mental health problems identified in the assessment.
- Encourage and provide influenza vaccination, regardless of the stage of pregnancy, during influenza season.

Routine Laboratory/diagnostic studies

- Blood type and screen.
- Complete Blood Count (CBC) for Hct/Hgb, MCV and Platelet Count.
- Hepatitis Screening: Hepatitis B surface antigen (HBsAg) and Hepatitis C antibody test.
- Syphilis screening.
- Screening for gestational diabetes if at high risk (see section on gestational diabetes below).
- HIV testing unless they decline (opt-out approach). For women that decline the provider should address objections and strongly encourage HIV screening.

Initial Approval Date and Reviews: New – November 2014. Nov 2016, Nov 2018, Nov 2020	Most Recent Revision and Approval Date: November 2020	Next Scheduled Review Date: November 2022 Condition: Perinatal Care
--	---	--

- Cervical Cancer Screening (if the patient is due).
- Urine C&S and urine dip for protein and glucose. Routine urine dip-stick testing is not recommended unless the woman has risk factors/symptoms of a urinary tract infection, renal disease, pre-eclampsia, unusual edema, or hypertension.

Genetic and infectious disease testing and counseling

Genetic Screening

- It is reasonable to offer Cystic fibrosis carrier screening to all couples regardless of ethnicity. Genetic counseling is recommended for individuals with a family history of cystic fibrosis or those found to be carriers.
- Hemoglobinopathy screening should be offered to individuals of African, Southeast Asian and Mediterranean descent. Couples at risk for having a child with sickle cell disease or thalassemia should be offered genetic counseling to review prenatal testing and reproduction options.
- Patients of Ashkenazi Jewish decent should be offered prenatal carrier screening for hereditary diseases common in this group.
- Screening for spinal muscular atrophy should be offered to all women who are considering pregnancy or are currently pregnant.⁴
- In patients with a family history of spinal muscular atrophy, molecular testing reports of the affected individual and carrier testing of the related parent should be reviewed, if possible, before testing. If the reports are not available, *SMNI* deletion testing should be recommended for the low-risk partner.⁴

Infectious Disease

- All pregnant women should be screened for chlamydia during the first prenatal visit. If positive, a test of cure should be offered to the patient four weeks after completing treatment and provide counseling to decrease risk of re-infection and refer partner for testing and treatment. Those women that are less than or equal to 25 years of age or at risk for chlamydia infection should be screened again during the third trimester.
- All pregnant women at risk for sexually transmitted diseases should be screened for gonorrhea at the initial prenatal visit. Risk factors include age less than 25, a previous infection, new or multiple sex partners, inconsistent condom use, commercial sex work and drug use. If positive, a test of cure should be offered to the patient four weeks after completing treatment and provide counseling to decrease risk of reinfection and refer partner for testing and treatment. Repeat screening is recommended during the third trimester of pregnancy.
- Rescreen for HIV in the third trimester for women at high risk of acquisition.
- Rescreen for syphilis in women at high risk of acquisition.
- All pregnant women should receive influenza immunization at any time of the year but especially from October through May. Trivalent or quadrivalent inactivated vaccines are all safe for mother and fetus at any point during the pregnancy. Live Attenuated vaccines should not be administered. There is no evidence of adverse consequences to thimerosal containing preparations and so they are safe to use.

Gestational Diabetes (GDM) Risk

Patients with the following risk factors should be screening for gestational diabetes at the first prenatal visit:

Pre-pregnancy BMI greater than or equal to 25 kg, and in Asian Americans, a BMI>23 and have one or more additional risk factors:

- Previous medical history of GDM.
- Physical Inactivity
- High risk race/ethnicity (ie. African American, Hispanic, Native American, Asian American, Pacific Islander)
- Women who have previously delivered a baby weighing 4000 g (~9lb) or more

<p>Initial Approval Date and Reviews: New – November 2014. Nov 2016, Nov 2018, Nov 2020</p>	<p>Most Recent Revision and Approval Date: November 2020</p>	<p>Next Scheduled Review Date: November 2022 Condition: Perinatal Care</p>
--	---	--

- Hypertension
- HDL cholesterol level <35mg/dL (0.90 mmol/L) and/or a triglyceride level >250 mg/DL (2.82 mmol/L)
- Women with polycystic ovarian syndrome
- HbA1c ≥5.7%, impaired fasting glucose, or impaired glucose tolerance
- Conditions associated with insulin resistance (i.e. pre-pregnancy BMI>40, acanthosis nigrans)
- First degree relative with DM

If gestational diabetes is not diagnosed, blood glucose testing should be repeated at 24-28 weeks of gestation

For Medicaid patients, complete and submit the Maryland Prenatal Risk Assessment form or the District of Columbia OB Risk form called the OB Global Authorization form at the time of the first prenatal visit.

Subsequent Prenatal Visits

Every visit

- Vital signs.
- Weight (height/weight/BMI – initial visit).
- Fetal assessment from 10th week.
- Uterine size for progressive growth and consistency with EDD.
- Domestic violence screening.
- Assessment of tobacco use and smoke exposure.
- Urine dip for protein and glucose if risk factors/symptoms of a urinary tract infection, renal disease, pre-eclampsia, unusual edema, or hypertension.

Specific Visit: 11 – 14 weeks

- Pelvic exam, if fetal heart tones (FHT) not heard with amplification.
- Breastfeeding has well documented short- and long-term medical and neurodevelopmental advantages for infants. As such, breastfeeding should be strongly encouraged during prenatal care as the best choice for feeding. Patients should be offered breastfeeding educational material and classes during pregnancy and provided resources for assistance after delivery.
- Review laboratory data. Offer iron supplementation for patients with anemia.
- Offer screening tests for aneuploidy, nuchal translucency sonogram and blood testing for First Trimester Screening. All pregnant women, regardless of age, should be counseled about non-invasive and invasive prenatal diagnostic testing for aneuploidy with a discussion of the risks and benefits of each. Women found to have increased risk for aneuploidy with non-invasive screening should be offered genetic counseling and the option of chorionic villus sampling (CVS) or second trimester amniocentesis.
- If previous low transverse cesarean delivery, discuss the risks, benefits, and alternatives to a trial of labor after cesarean as well as the risks and benefits of repeat cesarean delivery.

Specific Visit: 15-20 weeks

- Offer anatomic survey ultrasound to be completed at 18-20 weeks.
- Offer screening test for aneuploidy with a serum Multiple Marker Screen if the patient did not have first trimester screening (invasive or non-invasive) for aneuploidy. This also incorporates neural tube defect (NTD) screening. Screening and invasive diagnostic testing for aneuploidy should be available to all women who present for prenatal care before 20 weeks of gestation regardless of maternal age. Offer genetic counseling and the option of second trimester amniocentesis to women found to have increased risk for aneuploidy with screening.
- Offer neural tube defect screening (MSAFP) to women who elect first trimester screening or invasive testing for aneuploidy.
- Review signs and symptoms of pre-term labor (PTL).
- Review results of MSAFP/Multiple Marker screen and ultrasound if not already done.

<p><u>Initial Approval Date and Reviews:</u> New – November 2014, Nov 2016, Nov 2018, Nov 2020</p>	<p><u>Most Recent Revision and Approval Date:</u> November 2020</p>	<p><u>Next Scheduled Review Date:</u> November 2022 Condition: Perinatal Care</p>
---	--	---

Specific Visit: 24 – 28 weeks

- Screening for gestational diabetes. GTT if screen is abnormal.
- Select baby’s medical provider.
- Discuss normal fetal movement.
- Discuss prenatal classes.
- Discuss post-partum contraception. If applicable, patient should sign Medicaid consent for sterilization at this gestational age.

Specific Visit: 27-36 weeks

- TDAP (aka Tdap) should be administered during each pregnancy, irrespective of patient’s prior history of receiving. Optimal timing is between 27 and 36 weeks gestation to maximize maternal antibody response and passive antibody transfer levels in the newborn. Discuss with the patient that other adults who will be around her newborn, such as husbands, grandparents, older siblings, and babysitters, should also be vaccinated.

Specific Visit: 28 weeks

- Repeat type and screen if Rh negative, CBC for Hct/Hgb, MCV and platelet count
- Administer Rh-immune globulin if Rh (-) and indirect Coombs (-).
- Screen for gestational diabetes if not done at 24 wks.
- Discuss prenatal classes if not done at 24 wks.
- Confirm and document name of baby’s medical provider.
- Counsel woman about risks and benefits of circumcision.
- Discuss and encourage all women to breast feed.
- Discuss cord blood banking to allow a pregnant woman to make an informed decision on whether to participate in a public or private umbilical cord blood banking program. (Per PA House Bill 874).

Specific Visit: 32 – 34 weeks

- Repeat testing for women at risk for sexually transmitted disease, including syphilis screen, HIV, gonorrhea and chlamydia.
- Discuss Group B Strep screening and management protocol.
- Education and anticipatory guidance regarding breastfeeding.

Specific Visit: 36 weeks

- Determine fetal position.
- Group B Strep screen is performed at 35-37 weeks. Screening not needed if treatment in labor is indicated based on other risk factors such as group B strep bacteria during any trimester of the current pregnancy or previous infant with invasive GBS disease. Antibiotic resistance testing if Penicillin Allergic.
- Discuss the risks and benefits of HSV prophylaxis in women with a history of genital herpes.
- Labor education: latent phase of labor, rupture of membranes (ROM), active labor management, analgesia in labor.
- Counsel regarding labor induction indicating that in the absence of medical indications, labor should not be induced prior to 39 weeks gestation. Such early-term deliveries (37-38 6/7 weeks gestation) are associated with higher morbidity and mortality rates when compared to neonates and infants delivered between 39 weeks and 40 weeks of gestation.

Specific Visit: 38 weeks

- Review labor education; discuss again contraception, with an emphasis on the benefits of long-acting reversible contraception such as IUDs and implants.
- Education and anticipatory guidance regarding breastfeeding. Discuss patient’s preference: exclusive breastfeeding, mixed breastfeeding/formula, or formula.

Initial Approval Date and Reviews: New – November 2014. Nov 2016, Nov 2018, Nov 2020	Most Recent Revision and Approval Date: November 2020	Next Scheduled Review Date: November 2022 Condition: Perinatal Care
--	---	--

Specific Visit: > 41 weeks

- Baseline non-stress test (NST) or contraction stress test (CST), ultrasonography (US), biophysical profile (BPP) or a combination of these tests.
- Discuss labor induction > 41 weeks.

Postpartum Care

Postpartum vaccine

- Women (including women who are breastfeeding) who have not received a dose of Tdap previously should receive Tdap immediately after delivery and before discharge from the hospital. If Tdap cannot be administered before discharge, it should be administered as soon as feasible. Additionally, other family members and direct care caregivers should receive Tdap as recommended.

Postpartum visit

The most recent recommendations from ACOG encourage practitioners to consider the post partum care on ongoing process to address the health of the woman according to her individual needs. It is recommended that all women have contact with their OB provider within 3 weeks of delivery. Consider resources like telehealth or visiting health professionals for low resource areas or patients. This should be followed up with a comprehensive post partum visit no later than 12 weeks after birth. This is a change from the traditional 6-8 weeks and insurance carriers, including Medicaid, may not have aligned coverage with this recommendation,

Women with chronic conditions such as hypertension, obesity, diabetes, thyroid disorders, renal disease, and mood disorders should be counseled regarding timely follow up with OB or PCP. When a woman has a comorbidity like DM, HTN, mental health issues, etc. it is important to bring her back for a postpartum visit sooner than 6 weeks, more like 1-2 weeks, then 4-6 weeks, and even back again at 8 weeks. Have resources ready for your patients: internal medicine, psychiatry, cardiology, pelvic floor rehabilitation, etc.

A comprehensive postpartum visit should be completed no later than 12 weeks after birth. Many insurance carriers, especially Medicaid and the Medicaid Managed Care Organizations, still follow the traditional recommendation of on or between 21 days and 56 days after delivery.

Components of Postpartum Care

- Pelvic exam and /or weight, BP, breast, abdomen exam, wound check.
- Screen for postpartum depression and anxiety with a validated instrument. Refer for intervention if indicated.
- Follow up on preexisting mental health disorder, refer for or confirm attendance with appointments and titrate medications as appropriate for the post partum period.
- Screen for domestic violence.
- Screen for substance abuse and refer as appropriate.
- Screen for tobacco use and relapse and treat/refer as appropriate.
- Provide guidance regarding local services for mentoring and support.
- Discuss sexual activity, future family planning, and contraception with an emphasis on the benefits of long-acting reversible contraception and birth spacing.
- Review nutrition and exercise.
- Discuss method of infant feeding and the patient's infant feeding plan (exclusive breastfeeding for at least 6 months, formula, or mixed) and general infant care plan.
- Confirm infant has source of medical care.
- Discuss childcare strategy for return to work or school including immunizations for all care providers.
- Women with GDM should be screened for diabetes 6-12 weeks postpartum and should be followed up with subsequent screening for the development of diabetes or pre-diabetes.
- Evaluate for postpartum problems: perineal/C-section wound pain, urinary incontinence, fecal incontinence, dyspareunia/reduced sexual desire, fatigue/sleep issues, medication needs/titration

Initial Approval Date and Reviews: New – November 2014, Nov 2016, Nov 2018, Nov 2020	Most Recent Revision and Approval Date: November 2020	Next Scheduled Review Date: November 2022 Condition: Perinatal Care
--	---	--

- Confirm that delivery information is accurately documented, including complications, neonatal outcome.
- Perform health maintenance including immunization and Pap test and pelvic exam.
- Transition women to primary care or subspecialist health care providers for follow up care, as indicated.

<p><u>Initial Approval Date and Reviews:</u> New – November 2014. Nov 2016, Nov 2018, Nov 2020</p>	<p><u>Most Recent Revision and Approval Date:</u> November 2020</p>	<p><u>Next Scheduled Review Date:</u> November 2022 Condition: Perinatal Care</p>
---	--	---