

MFC Maryland Prior Authorization Table

Medication	FDA Indications* <i>*Note: Although every effort is made to keep this FDA indication list up to date, please consult the web link in the far-right column for the most accurate information.</i>	MFC Specifications	Manufacturer's Prescribing Info Link (Hold CTRL and click link to open)
Abecma (idecabtagene vicleucel)	Indicated for the treatment of adult patients with relapsed or refractory multiple myeloma after four or more prior lines of therapy, including an immunomodulatory agent, a proteasome inhibitor, and an anti-CD38 monoclonal antibody. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	ABECMA PI
Actimmune (interferon gamma-1b)	Indicated for: 1. reducing the frequency and severity of serious infections associated with Chronic Granulomatous Disease. 2. delaying time to disease progression in patients with severe, malignant osteopetrosis. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	ACTIMMUNE PI
Aimovig (erenumab-aooe)	Indicated for the preventive treatment of migraine in adults.	1. Rx by Neurologist 2. Member must have tried and failed at least 2 previous migraine prophylaxis medications. Examples of migraine prophylaxis medications include, but are not limited to, divalproex, metoprolol, propranolol, timolol, topiramate, amitriptyline, venlafaxine, atenolol.	AIMOVIG PI
Alecensa (alectinib)	Indicated for the treatment of patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.	Rx by Oncologist	ALECENSA PI
Alunbrig (brigatinib)	Indicated for the treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.	Rx by Oncologist	ALUNBRIG PI

Amitiza (lubiprostone)	Indicated for: 1. chronic idiopathic constipation (CIC) in adults. 2. opioid-induced constipation (OIC) in adult patients with chronic, non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation. Limitations of Use: Effectiveness of Amitiza in the treatment of OIC in patients taking diphenylheptane opioids (e.g., methadone) has not been established. 3. irritable bowel syndrome with constipation (IBS-C) in women ≥ 18 years old.	Member must have tried and failed at least 2 of the following: docusate, sennosides, polyethylene glycol 3350, lactulose, methylcellulose	AMITIZA PI
Amondys 45 (casimersen)	Indicated for the treatment of Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 45 skipping. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	AMONDYS 45 PI
Ampyra (dalfampridine)	Indicated to help improve walking in adults with MS.	1. Rx by Neurologist. 2. Documentation of MS with ambulatory dysfunction but must be able to walk 25 feet within 8-45 seconds at baseline. 3. Members must have a baseline gait assessment by PT within 90 days of beginning Ampyra. 4. Members must have a repeat evaluation after 3 months on Ampyra. Improvement in walking speed must be documented in order to obtain further refills. 5. Members must not have a history of seizure disorder or renal impairment.	AMPYRA PI
Austedo (deutetrabenazine)	Indicated for: 1. Chorea associated with Huntington’s disease. 2. Tardive dyskinesia in adults.		AUSTEDO PI
Ayvakit (avapritinib)	Indicated for the treatment of adults with unresectable or metastatic gastrointestinal stromal tumor harboring a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations.	Rx by Oncologist	AYVAKIT PI
Balversa (erdafitinib)	Indicated for the treatment of adult patients with locally advanced or metastatic urothelial carcinoma that has susceptible FGFR3 or FGFR2 genetic alterations and progressed during or following at least one line of prior platinum-containing chemotherapy including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.	Rx by Oncologist	BALVERSA PI

Benlysta (belimumab)	Indicated for the treatment of adult patients with active, autoantibody-positive, systemic lupus erythematosus who are receiving standard therapy.		BENLYSTA PI
Bethkis (tobramycin inh sol)	Indicated for management of cystic fibrosis patients with Pseudomonas aeruginosa.	Rx by Pulmonologist	BETHKIS PI
Blenprep (belantamab mafodotin-blmf)	Indicated for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least 4 prior therapies including an anti-CD38 monoclonal antibody, a proteasome inhibitor, and an immunomodulatory agent.	Rx by Oncologist	BLENPREP PI
Bosulif (bosutinib)	Indicated for the treatment of adult patients with: <ol style="list-style-type: none"> 1. newly diagnosed chronic phase Ph+ chronic myelogenous leukemia (CML). This indication is approved under accelerated approval based on molecular and cytogenetic response rates. Continued approval for this indication may be contingent upon verification and confirmation of clinical benefit in an ongoing long-term follow up trial. 2. chronic, accelerated, or blast phase Ph+ CML with resistance or intolerance to prior therapy. 	Rx by Oncologist	BOSULIF PI
Botox (onabotulinumtoxin A)	Indicated for: <ol style="list-style-type: none"> 1. treatment of overactive bladder (OAB) with symptoms of urge urinary incontinence, urgency, and frequency, in adults who have an inadequate response to or are intolerant of an anticholinergic medication. 2. treatment of urinary incontinence due to detrusor overactivity associated with a neurologic condition [e.g., spinal cord injury, multiple sclerosis] in adults who have an inadequate response to or are intolerant of an anticholinergic medication. 3. prophylaxis of headaches in adult patients with chronic migraine (≥ 15 days per month with headache lasting 4 hours a day or longer). 4. treatment of spasticity in patients 2 years of age and older. 5. treatment of cervical dystonia in adult patients, to reduce the severity of abnormal head position and neck pain. 6. treatment of severe axillary hyperhidrosis that is inadequately managed by topical agents in adult patients. 7. treatment of blepharospasm associated with dystonia in patients ≥ 12 years of age. 8. treatment of strabismus in patients ≥ 12 years of age. 	<ol style="list-style-type: none"> 1. Rx by Neurologist, Urologist, Ophthalmologist 2. Botox will NOT be approved for cosmetic purposes 	BOTOX PI
Breyanzi (lisocabtagene maraleucel)	Indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified (including DLBCL arising from indolent lymphoma), high-grade B-cell lymphoma, primary mediastinal large B-cell lymphoma, and follicular lymphoma grade 3B. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	BREYANZI PI

Braftovi (encorafenib)	Indicated: 1. in combination with binimetinib, for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDA-approved test. 2. in combination with cetuximab, for the treatment of adult patients with metastatic colorectal cancer (CRC) with a BRAF V600E mutation, as detected by an FDA-approved test, after prior therapy.	Rx by Oncologist	BRAFTOVI PI
Cabenuva (cabotegravir extended-release injectable suspension; rilpivirine extended-release injectable suspension)	Indicated as a complete regimen for the treatment of HIV-1 infection in adults to replace the current antiretroviral regimen in those who are virologically suppressed (HIV-1 RNA less than 50 copies per mL) on a stable antiretroviral regimen with no history of treatment failure and with no known or suspected resistance to either cabotegravir or rilpivirine.	Documentation of clinical appropriateness is required and MUST include the following: 1. most recent office note (<3 months old) with clear discussion of previous HIV regimen(s) and clinical response(s) to each. 2. lab test showing HIV-1 RNA less than 50 copies per mL (lab must be <3 months old). 3. evidence that the patient tolerated the oral lead-in with Vocabria.	CABENUVA PI
Cabometyx (cabozantinib)	Indicated for the treatment of: 1. patients with advanced renal cell carcinoma. 2. patients with hepatocellular carcinoma (HCC) who have been previously treated with sorafenib.	Rx by Oncologist	CABOMETYX PI
Cialis (tadalafil)	Indicated for the treatment of: 1. erectile dysfunction (ED) 2. the signs and symptoms of benign prostatic hyperplasia (BPH) 3. ED and the signs and symptoms of BPH (ED/BPH)	Rx by Pulmonologist or Cardiologist	CIALIS PI
Cinryze (C1 Esterase Inhibitor [Human])	Indicated for routine prophylaxis against angioedema attacks in adults, adolescents, and pediatric patients (6 years of age and older) with Hereditary Angioedema. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	CINRYZE PI
Cometriq (cabozantinib)	Indicated for treatment of progressive, metastatic medullary thyroid cancer.	Rx by Oncologist	COMETRIQ PI
Cosela (trilaciclib)	Indicated to decrease the incidence of chemotherapy-induced myelosuppression in adult patients when administered prior to a platinum/etoposide-containing regimen or topotecan-containing regimen for extensive-stage small cell lung cancer.	Rx by Oncologist	COSELA PI

Cotellic (cobimetinib)	Indicated for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, in combination with vemurafenib.	Rx by Oncologist	COTELLIC PI
Darzalex Faspro (daratumumab and hyaluronidase-fihj, SQ admin)	Indicated for the treatment of adult patients with multiple myeloma: <ol style="list-style-type: none"> in combination with bortezomib, melphalan and prednisone in newly diagnosed patients who are ineligible for autologous stem cell transplant in combination with lenalidomide and dexamethasone in newly diagnosed patients who are ineligible for autologous stem cell transplant and in patients with relapsed or refractory multiple myeloma who have received at least one prior therapy in combination with bortezomib and dexamethasone in patients who have received at least one prior therapy as monotherapy, in patients who have received at least three prior lines of therapy including a proteasome inhibitor (PI) and an immunomodulatory agent or who are double refractory to a PI and an immunomodulatory agent, 	Rx by Oncologist	DARZALEX FASPRO PI
DESMOPRESSIN NASAL SPRAY PRODUCTS: DDAVP spray- 0.01% Stimate spray- 1.5 mg/mL	DDAVP is indicated for: <ol style="list-style-type: none"> antidiuretic replacement therapy in the management of central cranial diabetes insipidus. treatment of transient polyuria and polydipsia post head trauma or surgery in the pituitary region. Stimate is indicated for: <ol style="list-style-type: none"> hemophilia A with Factor VIII coagulant activity levels greater than 5% - will stop bleeding in patients with hemophilia A with episodes of spontaneous or trauma-induced injuries such as hemarthroses, intramuscular hematomas or mucosal bleeding. mild to moderate classic von Willebrand’s disease (Type I) with Factor VIII levels greater than 5% - will stop bleeding in patients with episodes of spontaneous or trauma-induced injuries such as hemarthroses, intramuscular hematomas, mucosal bleeding or menorrhagia. 	STIMATE: Hemophilia A with factor VIII coagulant activity greater than 5%: <ul style="list-style-type: none"> ➤ *peri-operatively to prevent bleeding ➤ to treat spontaneous or trauma induced bleeding ***Note- Patients with factor VIII levels equal to or less than 5% or patients who have factor VIII antibodies are not candidates for the drug. It is contraindicated in patients under 3 months old. It is NOT indicated for Hemophilia B.	DDAVP FDA PI STIMATE PI
Descovy (emtricitabine and tenofovir alafenamide)	Indicated: <ol style="list-style-type: none"> in combination with other antiretroviral agents for the treatment of HIV- 1 infection in adults and pediatric patients weighing at least 35 kg. in combination with other antiretroviral agents other than protease inhibitors that require a CYP3A inhibitor for the treatment of HIV-1 infection in pediatric patients weighing at least 25 kg and less than 35 kg. 	Although Descovy is FDA approved for pre-exposure prophylaxis, MFC does not cover it for this indication. Descovy is covered for HIV treatment only. [MFC covers Truvada for pre-exposure prophylaxis].	DESCOVY PI

Dexcom G6 Continuous Glucose Monitoring (CGM) System	Indicated for the management of diabetes in persons age 2 years and older.	Rx by Endocrinologist. Please click link below for CGM Policy: MFC Continuous Glucose Monitoring Devices Policy	DEXCOM G6 PI
Dificid (fidaxomicin)	Indicated in adult and pediatric patients 6 months of age and older for the treatment of C. difficile-associated diarrhea.	Pt must have documented failures with both metronidazole and vancomycin, or contraindication(s) to the use of these agents.	DIFICID PI
Doptelet (avatrombopag)	Indicated for the treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure.	1. Rx by Hematologist 2. A recent (less than 1 month old) platelet count must be supplied with the clinical request, as well as information regarding the planned procedure.	DOPTELET PI
Dupixent (dupilumab)	Indicated: 1. for treatment of adult patients aged 12 years and older with moderate-to- severe atopic dermatitis whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable. DUPIXENT can be used with or without topical corticosteroids. 2. as add-on maintenance treatment in patients with moderate-to-severe asthma aged 12 years and older with an eosinophilic phenotype or with oral corticosteroid dependent asthma. 3. as add-on maintenance treatment in adult patients with inadequately controlled chronic rhinosinusitis with nasal polyposis.	Rx by Allergist or Dermatologist	DUPIXENT PI
Egrifta SV (tesamorelin injection)	Indicated for the reduction of excess abdominal fat in HIV-infected adult patients with lipodystrophy.		EGRIFTA SV PI
Eligard (leuprolide SQ)	see Leuprolide		
Elzonris (tagraxofusp-erzs)	Indicated for the treatment of blastic plasmacytoid dendritic cell neoplasm (BPDCN) in adults and in pediatric patients 2 years and older.	Rx by Oncologist	ELZONRIS PI
Empaveli (pegcetacoplan)	Indicated for the treatment of adult patients with paroxysmal nocturnal hemoglobinuria. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	EMPAVELI PI
Enhertu (fam-trastuzumab deruxtecan-nxk)	Indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received two or more prior anti-HER2- based regimens in the metastatic setting.	Rx by Oncologist	ENHERTU PI

<p>Epclusa (sofosbuvir/ velpatasvir)</p>	<p>EPCLUSA is a fixed-dose combination of sofosbuvir, a hepatitis C virus (HCV) nucleotide analog NS5B polymerase inhibitor, and velpatasvir, an HCV NS5A inhibitor, and is indicated for the treatment of patients with chronic HCV genotype (GT) 1, 2, 3, 4, 5 or 6 infection without cirrhosis and with compensated cirrhosis (Child-Pugh A) and with decompensated cirrhosis for use in combination with ribavirin (Child-Pugh B and C).</p> <p>EPCLUSA is also indicated for the treatment of adult and pediatric patients 6 years and older or weighing at least 17 kg with HCV genotypes 1, 2, 3, 4, 5, or 6 infection, who previously have been treated with a regimen containing an HCV NS5A inhibitor or an NS5B protease inhibitor, but not both.</p>	<p>For a full listing of all Prior Authorization requirements, please click the link below:</p> <p>Hepatitis C PRIOR AUTHORIZATION Submission Information</p>	<p>EPCLUSA PI</p>
<p>Erwinaze (asparaginase Erwinia chrysanthemi)</p>	<p>Indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of patients with ALL who have developed hypersensitivity to E. coli-derived asparaginase.</p>	<p>Rx by Oncologist</p>	<p>ERWINAZE PI</p>
<p>Esbriet (pirfenidone)</p>	<p>Indicated for the treatment of idiopathic pulmonary fibrosis IPF.</p>	<p>Rx by Pulmonologist or Cardiologist</p>	<p>ESBRIET PI</p>
<p>Evkeeza (evinacumab-dgnb)</p>	<p>Indicated as an adjunct to other low-density lipoprotein-cholesterol (LDL-C) lowering therapies for the treatment of adult and pediatric patients, aged 12 years and older, with homozygous familial hypercholesterolemia (HoFH).</p> <p>Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.</p>	<p>Must submit genetic testing confirming homozygous familial hypercholesterolemia (HoFH).</p> <p>***REQUIRES MFC PHYSICIAN REVIEW</p>	<p>EVKEEZA PI</p>
<p>Fasenra (benralizumab)</p>	<p>Indicated for the add-on maintenance treatment of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype.</p>	<p>Rx by Pulmonologist or Allergist</p>	<p>FASENRA PI</p>
<p>fentanyl</p>	<p>Indicated for the management of pain in opioid-tolerant patients, severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. Patients considered opioid- tolerant are those taking, for one week or longer, at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.</p>	<p>All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link:</p> <p>OPIOID PRIOR AUTHORIZATION FORM</p>	<p>fentanyl PI</p>
<p>Firazyr (icatibant)</p>	<p>Indicated for the treatment of acute attacks of hereditary angioedema (HAE) in adults ≥18 years of age (self-administered by the patient upon recognition of symptoms of an HAE attack after training under the guidance of a healthcare professional).</p>	<p>Rx by Allergist or ENT</p>	<p>FIRAZYR PI</p>
<p>Fotivda (tivozanib)</p>	<p>Indicated for the treatment of adult patients with relapsed or refractory advanced renal cell carcinoma (RCC) following two or more prior systemic therapies.</p>	<p>Rx by Hematology Oncology</p>	<p>FOTIVDA PI</p>

FreeStyle Libre Continuous Glucose Monitoring (CGM) System	Indicated for replacing blood glucose testing and detecting trends and tracking patterns aiding in the detection of episodes of hyperglycemia and hypoglycemia, facilitating both acute and long-term therapy adjustments in persons (age 18 and older) with diabetes.	Rx by Endocrinologist. Please click link below for CGM Policy: MFC Continuous Glucose Monitoring Devices Policy	FREESTYLE LIBRE CGM SYSTEM PI
Gavreto (pralsetinib)	Indicated for the treatment of adult patients with metastatic rearranged during transfection (RET) fusion- positive non-small cell lung cancer (NSCLC) as detected by an FDA approved test	Rx by Oncologist	GAVRETO PI
Gralise (gabapentin)	Indicated for the management of Postherpetic Neuralgia (PHN).		GRALISE PI
Growth Hormone	See Norditropin; See Serostim		
Haegarda (C1 Esterase Inhibitor SubQ (Human))	Indicated for routine prophylaxis to prevent Hereditary Angioedema attacks in patients 6 years of age and older.	Rx by Allergist/Immunologist	HAEGARDA PI
Hycamtin Capsules (topotecan)	Indicated for treatment of patients with relapsed small cell lung cancer.	Rx by Oncologist	HYCAMTIN PI
Ibrance (palbociclib)	Indicated for the treatment of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer in combination with: <ol style="list-style-type: none">1. an aromatase inhibitor as initial endocrine based therapy in postmenopausal women or in men;2. fulvestrant in patients with disease progression following endocrine therapy.	Rx by Oncologist	IBRANCE PI
Iclusig (ponatinib)	Indicated for: <ol style="list-style-type: none">1. treatment of adult patients with chronic phase, accelerated phase, or blast phase chronic myeloid leukemia or Ph+ ALL for whom no other tyrosine kinase inhibitor therapy is indicated.2. treatment of adult patients with T315I-positive CML (chronic phase, accelerated phase, or blast phase) or T315I-positive Philadelphia chromosome positive acute lymphoblastic leukemia.3. chromosome positive acute lymphoblastic leukemia.	Rx by Oncologist	ICLUSIG PI

<p>Imbruvica (ibrutinib)</p>	<p><u>Indicated for:</u></p> <ol style="list-style-type: none"> 1. Mantle cell lymphoma who have received at least one prior therapy. Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trials. 2. Chronic lymphocytic leukemia/small lymphocytic lymphoma. 3. Chronic lymphocytic leukemia/small lymphocytic lymphoma with 17p deletion. 4. Waldenström’s macroglobulinemia. 5. Marginal zone lymphoma who require systemic therapy and have received at least one prior anti-CD20-based therapy. Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial. 6. Chronic graft versus host disease after failure of one or more lines of systemic therapy. 	<p>Rx by Oncologist</p>	<p>IMBRUVICA PI</p>
<p>Insulin Pens</p>	<p>Age 0-18: Insulin pens will be provided for members ages 0-18 years of age.</p> <p>Age 19 and older: Basaglar and Tresiba pens do NOT require prior authorization and are available to all members. All other insulin pens require prior authorization.</p> <p>The MFC Insulin Pen Policy (Policy 211) states that insulin pens will be approved for members who cannot properly use and draw up insulin from vials into syringes or whose caregiver(s) cannot properly use and draw up insulin from vials into syringes including members with:</p> <ol style="list-style-type: none"> 1) Poor visual acuity, 2) Poor manual dexterity, or 3) Limited ability to learn proper technique due to educational challenges <p>Medical records may be required to support the PA request.</p>		
<p>Jakafi (ruxolitinib)</p>	<p><u>Indicated for:</u></p> <ol style="list-style-type: none"> 1. treatment of patients with intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis, and post- essential thrombocythemia myelofibrosis in adults. 2. treatment of adults with polycythemia vera who have had an inadequate response to or are intolerant of hydroxyurea. 3. treatment of steroid-refractory acute graft-versus-host disease in adult and pediatric patients 12 years and older. 	<p>Rx by Hematologist/Oncologist</p>	<p>JAKAFI PI</p>

Jivi (empagliflozin)	Recombinant DNA-derived, Factor VIII concentrate indicated for use in previously treated adults and adolescents (12 years of age and older) with hemophilia A (congenital Factor VIII deficiency) for: <ol style="list-style-type: none"> 1. On-demand treatment and control of bleeding episodes. 2. Perioperative management of bleeding. 3. Routine prophylaxis to reduce the frequency of bleeding episodes. 	Rx by Hematologist	JIVI PI
Juxtapid (lomitapide)	Indicated as an adjunct to a low-fat diet and other lipid-lowering treatments, including LDL apheresis where available, to reduce LDL-C, total cholesterol, apolipoprotein B, and non-HDL-C in patients with homozygous familial hypercholesterolemia.	Rx by Cardiology or Endocrinologist	JUXTAPID PI
Jynarque (tolvaptan)	Indicated to slow kidney function decline in adults at risk of rapidly progressing autosomal dominant polycystic kidney disease.	Rx by Nephrologist	JYNARQUE PI
Kalbitor (ecallantide)	Indicated for treatment of acute attacks of hereditary angioedema (HAE) in patients 12 years of age and older.	Rx by Immunologist or Allergist	KALBITOR PI
Kalydeco (ivacaftor)	Indicated for the treatment of cystic fibrosis (CF) in patients age 6 months and older who have one mutation in the CFTR gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient’s genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.	Rx by Pulmonologist	KALYDECO PI
Kisqali (ribociclib)	Indicated in combination with: <ol style="list-style-type: none"> 1. an aromatase inhibitor for the treatment of pre/perimenopausal or postmenopausal women with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer, as initial endocrine-based therapy; or 2. fulvestrant for the treatment of postmenopausal women with HER2-positive, HER2-negative advanced or metastatic breast cancer, as initial endocrine based therapy or following disease progression on endocrine therapy. 	Rx by Oncologist	KISQALI PI

<p>Kymriah (tisagenlecleucel)</p>	<p>Indicated for:</p> <ol style="list-style-type: none"> 1. Patients up to 25 years of age with B-cell precursor acute lymphoblastic leukemia (ALL) that is refractory or in second or later relapse. 2. Adult patients with relapsed or refractory (r/r) large B-cell lymphoma after two or more lines of systemic therapy including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, high grade B-cell lymphoma and DLBCL arising from follicular lymphoma. <p>In accordance with criteria developed by the Maryland Medicaid Program, MedStar Family Choice considers Kymriah medically necessary when all of the following conditions are met:</p> <ol style="list-style-type: none"> a. Recipient has relapsed or refractory B-cell ALL, defined as Second or greater bone marrow relapse; OR b. Any bone marrow relapse after allogeneic stem cell transplantation; OR c. Primary refractory as defined by not achieving a complete remission after 2 cycles of a standard chemotherapy regimen or chemorefractory as defined by not achieving a complete remission after 1 cycle of standard chemotherapy for relapsed leukemia; OR d. Patients with Philadelphia chromosome positive (Ph+) ALL are eligible if they are intolerant to or have failed 2 lines of tyrosine kinase inhibitor therapy (TKI), or if TKI therapy is contraindicated; AND e. Recipient is 25 years of age or younger; AND f. Documentation of CD19 tumor expression; AND g. Performance score on Karnofsky or Lansky Scale is greater than or equal to 50%; AND h. Life expectancy > 12 weeks; AND i. Adequate cardiac, pulmonary, and other organ function (as determined by protocol from treatment facility); AND j. The treatment facility that dispenses and administers Kymriah is enrolled and complies with the Risk Evaluation and Mitigation Strategy; AND k. One-time, single administration with dosing in accordance with the FDA label. <p>Kymriah is considered investigational and not medically necessary when the above medically necessary criteria are not met, and for all other indications, including but not limited to:</p> <ol style="list-style-type: none"> 1. Isolated extra-medullary disease relapse; or 2. Patients with Burkitt's lymphoma/leukemia (i.e. patients with mature B-cell ALL, leukemia with B-cell [sIg positive and kappa or lambda restricted positivity] ALL, with FAB L3 morphology and /or a MYC translocation); or 3. Prior malignancy, except carcinoma in situ of the skin or cervix treated with curative intent and with no evidence of active disease; or 4. Treatment with any other chimeric antigen receptor therapy or genetically modified T cell therapy; or 	<p>Rx by Oncologist, see criteria to the left.</p>	<p>KYMRIAH PI</p>
--	--	--	-----------------------------------

	<ol style="list-style-type: none"> 5. Any active uncontrolled infection; or 6. Hepatitis B or C (if viral load is detectable); or 7. Human Immunodeficiency Virus (HIV); or 8. Presence of grade 2 to 4 acute or extensive chronic graft-versus-host disease (GVHD); or 9. Active CNS involvement by malignancy, defined by CNS-3 per NCCN guidelines. 		
<p>LEUPROLIDE PRODUCTS:</p> <p>Eligard (leuprolide SQ)</p> <p>Lupron (leuprolide acetate)</p> <p>Lupron Depot (leuprolide acetate for depot suspension)</p> <p>Lupron Depot-PED (leuprolide acetate for depot suspension)</p>	<p>Indicated for:</p> <ol style="list-style-type: none"> 1. palliative treatment for advanced prostate cancer (Eligard). 2. treatment of pediatric patients with central precocious puberty (Lupron Depot- PED). 3. treatment of endometriosis (Lupron and Lupron Depot). 4. Management of endometriosis, including pain relief, recurrence symptoms and reduction of endometriotic lesions, (Lupron and Lupron Depot). 5. uterine leiomyomata (fibroids) along with concurrent iron therapy in preparation for surgery [duration of treatment should be for 6 months or less (Lupron and Lupron Depot)]. 		<p>ELIGARD PI</p> <p>LUPRON 3.75 mg PI</p> <p>LUPRON DEPOT 11.25 MG PI</p> <p>LUPRON DEPOT- PED PI</p>
<p>Libtayo (cemiplimab-rwlc)</p>	<p>Indicated for the treatment of patients with metastatic cutaneous squamous cell carcinoma (CSCC) or locally advanced CSCC who are not candidates for curative surgery or curative radiation.</p>	Rx by Oncologist	LIBTAYO PI
<p>LO Loestrin Fe (norethindrone, ethinyl estradiol and ferrous fumarate)</p>	<p>See Oral Contraceptive</p>	<p>ANY OCP on prior authorization requires documentation demonstrating a compelling reason why formulary OCPs cannot be used [ex: intolerance, prior side effects, failures, etc. documented after a 3-month trial of formulary OCPs]</p>	LO LOESTRIN PI
<p>Lorbrena (lorlatinib)</p>	<p>Indicated for the treatment of patients with anaplastic lymphoma kinase (ALK)- positive metastatic non-small cell lung cancer (NSCLC) whose disease has progressed on:</p> <ul style="list-style-type: none"> • crizotinib and at least one other ALK inhibitor for metastatic disease; or • alectinib as the first ALK inhibitor therapy for metastatic disease; or • ceritinib as the first ALK inhibitor therapy for metastatic disease 	Rx by Oncologist	LORBRENA PI

Lovaza (omega-3-acid ethyl esters) (historical Omacor)	Indicated as an adjunct to diet to reduce triglyceride levels in adult patients with <ul style="list-style-type: none"> severe (≥ 500 mg/dL) hypertriglyceridemia. 		LOVAZA PI
Lumakras (sotorasib)	Indicated for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an FDA-approved test, who have received at least one prior systemic therapy.	Rx by Oncologist	LUMAKRAS PI
Lumoxiti (moxetumomab pasudotox-tdfk)	Indicated for the treatment of adult patients with relapsed or refractory hairy cell leukemia (HCL) who received at least 2 prior systemic therapies, including treatment with a purine nucleoside analog.	Rx by Oncologist	LUMOXITI PI
Lupron and Lupron Depot	See Leuprolide		
Lupkynis (voclosporin)	Indicated in combination with a background immunosuppressive therapy regimen for the treatment of adult patients with active lupus nephritis.		LUPKYNIS PI
Lynparza (olaparib)	<u>Indicated for:</u> <ol style="list-style-type: none"> First-Line Maintenance Treatment of BRCA-mutated Advanced Ovarian Cancer - For the maintenance treatment of adult patients with deleterious or suspected deleterious germline or somatic <i>BRCA</i>-mutated (<i>gBRCAm</i> or <i>sBRCAm</i>) advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for Lynparza. In combination with bevacizumab for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube or primary peritoneal cancer who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD) positive status defined by either: a deleterious or suspected deleterious BRCA mutation, and/or genomic instability. Maintenance Treatment of Recurrent Ovarian Cancer- For the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, who are in complete or partial response to platinum-based chemotherapy. Advanced <i>gBRCA</i>-mutated Ovarian Cancer After 3 or More Lines of Chemotherapy- For the treatment of adult patients with deleterious or suspected deleterious germline <i>BRCA</i>-mutated (<i>gBRCAm</i>) advanced ovarian cancer who have been treated with 3 or more prior lines of chemotherapy. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA. 	Rx by Oncologist	LYNPARZA PI

	<p>5. Germline BRCA-mutated HER2-negative Metastatic Breast Cancer – In patients with deleterious or suspected deleterious <i>gBRCAm</i>, human epidermal growth factor receptor 2 (HER2)-negative metastatic breast cancer who have been treated with chemotherapy in the neoadjuvant, adjuvant or metastatic setting. Patients with hormone receptor (HR)- positive breast cancer should have been treated with a prior endocrine therapy or be considered inappropriate for endocrine therapy. Select patients for therapy based on FDA-approved companion diagnostic for Lynparza.</p> <p>6. First-Line Maintenance <i>gBRCAm</i> Metastatic Pancreatic Cancer-- For the maintenance treatment of adult patients with deleterious or suspected deleterious <i>gBRCAm</i> metastatic pancreatic adenocarcinoma whose disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.</p> <p>7. HRR Gene-mutated Metastatic Castration-Resistant Prostate Cancer For the treatment of adult patients with deleterious or suspected deleterious germline or somatic homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer (mCRPC) who have progressed following prior treatment with enzalutamide or abiraterone. Select patients for therapy based on an FDA-approved companion diagnostic for LYNPARZA.</p>		
Macrilen (macimorelin)	Indicated for the diagnosis of adult growth hormone deficiency.	Rx by Endocrinologist	MACRILEN PI
Mavyret (glecaprevir and pibrentasvir)	MAVYRET is a fixed-dose combination of glecaprevir, a hepatitis C virus (HCV) NS3/4A protease inhibitor, and pibrentasvir, an HCV NS5A inhibitor, and is indicated for the treatment of patients with chronic HCV genotype (GT) 1, 2, 3, 4, 5 or 6 infection without cirrhosis and with compensated cirrhosis (Child-Pugh A). MAVYRET is also indicated for the treatment of adult and pediatric patients 12 years and older or weighing at least 45 kg with HCV genotype 1 infection, who previously have been treated with a regimen containing an HCV NS5A inhibitor or an NS3/4A protease inhibitor, but not both.	For a full listing of all Prior Authorization requirements, please click the link below: Hepatitis C PRIOR AUTHORIZATION Submission Information	MAVYRET PI
Mekinist (trametinib)	Indicated: <ol style="list-style-type: none"> 1. BRAF V600E or V600K Mutation-Positive Unresectable or Metastatic Melanoma - as a single agent in BRAF-inhibitor treatment-naïve patients or in combination with dabrafenib, for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E or V600K mutations, as detected by an FDA-approved test. 2. Adjuvant Treatment of BRAF V600E or V600K Mutation-Positive Melanoma- indicated in combination with dabrafenib, for the adjuvant treatment of patients with melanoma with BRAF V600E or V600K mutations as detected by an FDA-approved test, and involvement of lymph node(s), following complete resection. 3. BRAF V600E Mutation-Positive Metastatic NSCLC- in combination with dabrafenib, for the treatment of patients with metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation as detected by an FDA-approved test. 	Rx by Oncologist	MEKINIST PI

	4. BRAF V600EMutation-Positive Locally Advanced or Metastatic Anaplastic Thyroid Cancer - in combination with dabrafenib, for the treatment of patients with locally advanced or metastatic anaplastic thyroid cancer.		
Mektovi (binimetinib)	Indicated, in combination with encorafenib, for the treatment of patients with unresectable or metastatic melanoma with a BRAF V600E or V600K mutation, as detected by an FDA-approved test.	Rx by Oncologist	MEKTOVI PI
methadone (for pain)	Indicated for management of pain severe enough to require daily, around-the- clock, long-term opioid treatment and for which alternative treatment options are inadequate.	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTHORIZATION FORM	METHADONE PI
Minastrin 24 Fe (norethindrone, ethinyl estradiol and ferrous fumarate)	See Oral Contraceptive ***CHEWABLE	ANY OCP on prior authorization requires documentation demonstrating a compelling reason why formulary OCPs cannot be used [ex: intolerance, prior side effects, failures, etc. documented after a 3-month trial of formulary OCPs]	MINASTRIN 24 Fe PI
Movantik (naloxegol)	Indicated for the treatment of opioid-induced constipation in adults with chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation.	Failure of at least 2 of the following: docusate, sennosides, polyethylene glycol 3350, lactulose, methylcellulose	MOVANTIK PI
MS Contin (morphine sulfate controlled release)	Indicated for the management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period.	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTHORIZATION FORM	MS CONTIN PI
Mulpleta (lusutrombopag)	Indicated for the treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure.		MULPLETA PI
Natazia (estradiol valerate and estradiol valerate/dienogest)	Indicated for: 1. use by women to prevent pregnancy. 2. treatment of heavy menstrual bleeding in women without organic pathology who choose to use an oral contraceptive as their method of contraception.	ANY OCP on prior authorization requires documentation demonstrating a compelling reason why formulary OCPs cannot be used [ex: intolerance, prior side effects, failures, etc. documented after a 3-month trial of formulary OCP(s)]	NATAZIA PI
Norditropin (somatropin (fDNA origin) injection)	Indicated for: 1. Pediatric: Treatment of pediatric patients with growth failure due to inadequate secretion of endogenous growth hormone (GH), short stature associated with Noonan syndrome, short stature associated with Turner syndrome, short stature born small for gestational age (SGA) with no catch- up growth by age 2 to 4 years, Idiopathic Short Stature (ISS), and growth failure due to Prader-Willi Syndrome.	Rx by Endocrinologist	NORDITROPIN PI

	2. Adult: Replacement of endogenous GH in adults with growth hormone deficiency.		
Nourianz (istradefylline)	Indicated as adjunctive treatment to levodopa/carbidopa in adult patients with Parkinson's disease experiencing "off" episodes.	Rx by Neurologist	NOURIANZ PI
NovoSeven RT (Coagulation Factor VIIa recombinant)	Indicated for: <ol style="list-style-type: none"> treatment of bleeding episodes and perioperative management in adults and children with hemophilia A or B with inhibitors, congenital Factor VII (FVII) deficiency, and Glanzmann's thrombasthenia with refractoriness to platelet transfusions, with or without antibodies to platelets. treatment of bleeding episodes and perioperative management in adults with acquired hemophilia. <p>Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.</p>	***REQUIRES MFC PHYSICIAN REVIEW	NOVOSEVEN RT PI
Noxafil (posaconazole)	Indicated for: <ol style="list-style-type: none"> prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised, such as HSCT recipients with GVHD or those with hematologic malignancies with prolonged neutropenia from chemotherapy. treatment of oropharyngeal candidiasis (OPC), including OPC refractory (rOPC) to itraconazole and/or fluconazole. 	Rx by ID	NOXAFIL PI
Nubeqa (darolutamide)	Indicated for the treatment of patients with non-metastatic castration-resistant prostate cancer.	Rx by Oncologist or Urologist	NUBEQA PI
Nucala (mepolizumab)	Indicated for: <ol style="list-style-type: none"> The add-on maintenance treatment of patients with severe asthma aged 6 years and older, and with an eosinophilic phenotype. The treatment of adult patients with eosinophilic granulomatosis with polyangiitis. The treatment of adult and pediatric patients aged 12 years and older with hypereosinophilic syndrome for ≥6 months without an identifiable non-hematologic secondary cause. 	Rx by Allergist or Pulmonologist	NUCALA PI
Nulibry (fosdenopterin)	Indicated to reduce the risk of mortality in patients with molybdenum cofactor deficiency (MoCD) Type A. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	NULIBRY PI
Ofev (nintedanib)	Indicated for: <ol style="list-style-type: none"> The treatment of idiopathic pulmonary fibrosis. Treatment of chronic fibrosing interstitial lung diseases with a progressive 	Rx by Pulmonologist	OFEV PI

	phenotype. 3. To slow the rate of decline in pulmonary function in patients with systemic sclerosis associated interstitial lung disease.		
OmniPod-Insulin Management (EIM) Systems	Indicated for subcutaneous delivery of insulin at set and variable rates for the management of diabetes mellitus in persons requiring insulin and for the quantitative measurement of glucose in fresh whole capillary blood (in vitro) from the finger.	RX Endocrinology Must meet criteria found in Policy 1413 Please click link below for EIM Policy: MFC External Insulin Pumps Policy	OMNIPOD PI
Onpattro (patisiran)	Indicated for the treatment of the polyneuropathy of hereditary transthyretin-mediated amyloidosis in adults.	Rx by Rheumatology or Neurology	ONPATTRO PI
Onureg (azacitidine)	Indicated for continued treatment of adult patients with acute myeloid leukemia who achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy and are not able to complete intensive curative therapy.	Rx by Oncologist	ONUREG PI
OPIOIDS PRIOR AUTHORIZATION TERMS	FOR IMPORTANT INFORMATION ABOUT PRESCRIBING OPIOIDS FOR MEDSTAR FAMILY CHOICE MEMBERS, PLEASE CLICK THE LINK BELOW: https://www.medstarfamilychoice.com/maryland-healthchoice/for-maryland-healthchoice-physicians/pharmacy/opioid/	The Opioid PA form can be accessed using the following link: OPIOID PRIOR AUTHORIZATION FORM	
Oral Contraceptives	While some oral contraceptives have additional indications (ex: Beyaz for acne, PMDD, folate replacement; Estrostep Fe for acne; Safyral for folate replacement; Natazia for heavy periods), most are simply indicated for the prevention of pregnancy.	ANY OCP on prior authorization requires documentation demonstrating a compelling reason why formulary OCPs cannot be used [ex: intolerance, prior side effects, failures, etc. documented after a 3-month trial of formulary OCPs]	
Orfadin (nitisinone)	Indicated for the treatment of adult and pediatric patients with hereditary tyrosinemia type 1 in combination with dietary restriction of tyrosine and phenylalanine. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	ORFADIN PI
Orkambi (lumacaftor/ivacaftor)	Indicated for the treatment of cystic fibrosis in patients age 2 years and older who are homozygous for the F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene.	Rx by Pulmonologist	ORKAMBI PI
Oriahnn (elagolix, estradiol, and norethindrone acetate capsules)	Indicated for the management of heavy menstrual bleeding associated with uterine leiomyomas in premenopausal women.	Must have tried and failed hormonal options (OCP, progesterone, hormonal IUD) or have a contraindication to using these therapies	ORIAHNN PI

Orilissa (elagolix)	Indicated for the management of moderate to severe pain associated with endometriosis.		ORLISSA PI
Orladeyo (berotralstat)	Indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adults and pediatric patients 12 years and older. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	ORLADEYO PI
Oxlumo (lumasiran)	Indicated for the treatment of primary hyperoxaluria type 1 (PH1) to lower urinary oxalate levels in pediatric and adult patients. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	OXLUMO PI
Oxymorphone ER	Indicated for the relief of moderate to severe pain in patients requiring continuous around-the-clock opioid treatment for an extended period of time.	All long-acting opioids require Prior Authorization (PA). The PA form can be accessed using the following link: OPIOID PRIOR AUTHORIZATION FORM	OPANA ER PI
Pemazyre (pemigatinib)	Indicated for the treatment of adults with previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with a fibroblast growth factor receptor 2 fusion or other rearrangement as detected by an FDA-approved test.	Rx by Oncologist	PEMAZYRE PI
Pretomanid	Indicated, as part of a combination regimen with bedaquiline and linezolid for the treatment of adults with pulmonary extensively drug resistant, treatment-intolerant or nonresponsive multidrug-resistant tuberculosis.	Rx by Pulmonologist	PRETOMANID
Piqray (alpelisib)	Indicated in combination with fulvestrant for the treatment of postmenopausal women, and men, with hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated, advanced or metastatic breast cancer as detected by an FDA-approved test following progression on or after an endocrine-based regimen.	Rx by Oncologist	PIQRAY PI
Polivy (polatuzumab vedotin)	Indicated in combination with bendamustine and a rituximab product for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma, not otherwise specified, after at least two prior therapies.	Rx by Oncologist	POLIVY PI
Prolia (denosumab)	Indicated for: 1. treatment of postmenopausal women with osteoporosis at high risk for fracture. 2. treatment to increase bone mass in men with osteoporosis at high risk for fracture. 3. treatment of glucocorticoid-induced osteoporosis in men and women at high risk for fracture. 4. treatment to increase bone mass in men at high risk for fracture receiving		PROLIA PI

	androgen deprivation therapy for non-metastatic prostate cancer. 5. treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer.		
Pulmozyme (dornase alfa) Inhalation solution	Indicated in conjunction with standard therapies for the management of cystic fibrosis patients to improve pulmonary function.	Rx by Pulmonologist	PULMOZYME PI
Qbrexza (glycopyrronium)	Indicated for topical treatment of primary axillary hyperhidrosis in adults and pediatric patients 9 years of age and older.	1. Must have tried and failed OTC Clinical Strength antiperspirants and at least one prescription strength antiperspirant (ex: Drysol). 2. Documentation that symptoms are persistent despite previous treatment attempts and that the degree of symptomatology impacts quality of life must be clearly indicated in a recent (<6 month old) clinical encounter note.	QBREXZA PI
Rasuvo (methotrexate inj)	Indicated for: 1. Management of patients with severe, active rheumatoid arthritis and polyarticular juvenile idiopathic arthritis, who are intolerant of or had an inadequate response to first-line therapy. 2. Symptomatic control of severe, recalcitrant, disabling psoriasis in adults who are not adequately responsive to other forms of therapy.	Rx by Rheumatology or Dermatology	RASUVO PI
Ravicti (glycerol phenylbutyrate)	Indicated for chronic management of patients with urea cycle disorders (UCDs) who cannot be managed by dietary protein restriction and/or amino acid supplementation alone. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	RAVICTI PI
Repatha (evolocumab)	Indicated: 1. to reduce the risk of myocardial infarction, stroke, and coronary revascularization in adults with established cardiovascular disease. 2. as an adjunct to diet, alone or in combination with other lipid-lowering therapies (e.g., statins, ezetimibe), for treatment of adults with primary hyperlipidemia (including heterozygous familial hypercholesterolemia) to reduce low-density lipoprotein cholesterol. 3. as an adjunct to diet and other LDL-lowering therapies (e.g., statins, ezetimibe, LDL apheresis) in patients with homozygous familial hypercholesterolemia who require additional lowering of LDL-C.	Rx by Cardiologist or Lipid Specialist.	REPATHA PI
Restasis (cyclosporine ophthalmic emulsion)	Indicated to increase tear production in patients whose tear production is presumed to be suppressed due to ocular inflammation associated with keratoconjunctivitis sicca.		RESTASIS PI

Retevmo (selpercatinib)	Indicated for: 1. Adult patients with metastatic RET (rearranged during transfection) fusion-positive non-small cell lung cancer (NSCLC). 2. Adult and pediatric patients 12 years of age and older with advanced or metastatic RET-mutant medullary thyroid cancer (MTC) who require systemic therapy. 3. Adult and pediatric patients 12 years of age and older with advanced or metastatic RET fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if radioactive iodine is appropriate).	Rx by Oncologist	RETEVMO PI
Revatio (sildenafil)	Indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group I) in adults to improve exercise ability and delay clinical worsening. Studies establishing effectiveness were short-term (12 to 16 weeks) and included predominately patients with NYHA Functional Class II–III symptoms. Etiologies were idiopathic (71%) or associated with connective tissue disease (25%).	Rx by Pulmonologist or Cardiologist	REVATIO PI
Revcovi (elapegedemase-lvlr)	Indicated for the treatment of adenosine deaminase severe combined immune deficiency (ADA-SCID) in pediatric and adult patients. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	REVCОВI PI
Revlimid (lenalidomide)	<u>Indicated for the treatment of adult patients with:</u> 1. Multiple myeloma (MM), in combination with dexamethasone. 2. MM, as maintenance following autologous hematopoietic stem cell transplantation (auto-HSCT). 3. Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndromes. 4. (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities. 5. Mantle cell lymphoma (MCL) whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib. 6. Previously treated follicular lymphoma (FL), in combination with a rituximab product. 7. Previously treated marginal zone lymphoma (MZL), in combination with a rituximab product		REVLIMID PI
Reyvow (lasmiditan)	Indicated for the acute treatment of migraine with or without aura in adults.	Member must have tried and failed NSAIDs and Triptans or have a contraindication to taking either of these medications.	REYVOW PI
Rituxan Hycela (rituximab and hyaluronidase human)	Indicated for: 1. Follicular Lymphoma (FL): <ul style="list-style-type: none"> • Relapsed or refractory, follicular lymphoma as a single agent. • Previously untreated follicular lymphoma in combination with first line chemotherapy and, in patients achieving a complete or partial response to 	Rx by Oncologist	RITUXAN HYCELA PI

	<p>rituximab in combination with chemotherapy, as single agent maintenance therapy.</p> <ul style="list-style-type: none"> • Non-progressing (including stable disease), follicular lymphoma as a single agent after first-line cyclophosphamide, vincristine, and prednisone (CVP) chemotherapy. <p>2. Diffuse Large B-cell Lymphoma (DLBCL) previously untreated diffuse large B-cell lymphoma in combination with cyclophosphamide, doxorubicin, vincristine, prednisone (CHOP) or other anthracycline-based chemotherapy regimens.</p> <p>3. Chronic Lymphocytic Leukemia (CLL) previously untreated and previously treated CLL in combination with fludarabine and cyclophosphamide (FC).</p>		
Rozlytrek (entrectinib)	<p>Indicated for the treatment of:</p> <ol style="list-style-type: none"> 1. Adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive. 2. Adult and pediatric patients 12 years of age and older with solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity and have either progressed following treatment or have no satisfactory alternative therapy. 	Rx by Oncologist	ROZLYTREK
Rubraca (rucaparib)	<p>Indicated for:</p> <ol style="list-style-type: none"> 1. maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy. 2. treatment of adult patients with deleterious BRCA mutation (germline and/or somatic)-associated epithelial ovarian, fallopian tube, or primary peritoneal cancer who have been treated with two or more chemotherapies. 	Rx by Oncologist	RUBRACA PI
Rybrevant (amivantamab-vmjw)	<p>Indicated for the treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 20 insertion mutations, as detected by an FDA-approved test, whose disease has progressed on or after platinum-based chemotherapy.</p>	Rx by Oncologist	RYBREVANT PI
Santyl Ointment Collagenase	<p>Indicated for debriding chronic dermal ulcers and severely burned areas.</p>	Rx by Dermatologist or Wound Care Specialist	SANTYL PI
Serostim (somatropin (rDNA origin))	<p>Indicated for the treatment of HIV patients with wasting or cachexia to increase lean body mass and body weight and improve physical endurance.</p>	Rx by ID or HIV Specialist	SEROSTIM PI
Seysara (seracycline)	<p>Indicated for the treatment of inflammatory lesions of non-nodular moderate to severe acne vulgaris in patients 9 years of age and older.</p>	<ol style="list-style-type: none"> 1. Rx by Dermatologist. 2. Failure of at least one other oral tetracycline antibiotic. 	SEYSERA PI
Signifor LAR (pasireotide)	<p>Indicated for:</p> <ol style="list-style-type: none"> 1. Patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option. 	Rx by Endocrinologist	SIGNIFOR LAR PI

	2. Patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative.		
Sirturo (bedaquiline)	Indicated as part of combination therapy in adult and pediatric patients (5 years and older and weighing at least 15 kg) with pulmonary multi-drug resistant tuberculosis (MDR-TB). Reserve SIRTURO for use when an effective treatment regimen cannot otherwise be provided.	Rx by ID	SITURO PI
Soliris (eculizumab)	Indicated for: 1. treatment of patients with paroxysmal nocturnal hemoglobinuria (PNH) to reduce hemolysis. 2. treatment of patients with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy. The effectiveness of Soliris in aHUS is based on the effects on thrombotic microangiopathy (TMA) and renal function. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	SOLIRIS PI
Spinraza (nusinersen)	Indicated for the treatment of spinal muscular atrophy (SMA) in pediatric and adult patients. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	SPINRAZA PI
Stimate nasal spray (desmopressin)	See Desmopressin Products		
Stivarga (regorafenib)	Indicated for: 1. treatment of metastatic colorectal cancer previously treated with ALL the following therapies: a. fluoropyrimidine-based chemotherapy b. oxaliplatin-based chemotherapy c. irinotecan-based chemotherapy d. an anti-vascular endothelial growth factor (VEGF) therapy e. if Kirsten RNA Associated Rat Sarcoma 2 Virus Gene (KRAS) wild type, an anti-epidermal growth factor receptor (EGFR) therapy 2. treatment of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST), previously treated with imatinib mesylate and sunitinib malate. 3. hepatocellular carcinoma (HCC) who have been previously treated with sorafenib.	Rx by Oncologist	STIVARGA PI
Stromectol (ivermectin)	Indicated for the treatment of: 1. Strongyloidiasis of the intestinal tract (i.e., nondisseminated) strongyloidiasis due to Strongyloides stercoralis. 2. Onchocerciasis due to the nematode parasite Onchocerca volvulus.	At this time, outpatient use for COVID-19 treatment is prohibited.	STROMECTOL PI

Synagis (palivizumab)	<p>Indicated for prevention of serious lower respiratory tract disease caused by RSV in pediatric Patients:</p> <ol style="list-style-type: none"> with a history of premature birth (≤ 35 weeks gestational age) and who are 6 months of age or younger at the beginning of RSV season with bronchopulmonary dysplasia (BPD) that required medical treatment within the previous 6 months and who are 24 months of age or younger at the beginning of RSV season with hemodynamically significant congenital heart disease (CHD) and who are 24 months of age or younger at the beginning of RSV season. <p>MedStar Family Choice uses the newest recommendations of the American Academy of Pediatrics (AAP). Recommendations were last updated in the journal Pediatrics (7/28/2014 issue): Updated Guidance for Palivizumab Prophylaxis Among Infants and Young Children at Increased Risk of Hospitalization for Respiratory Syncytial Virus Infection.</p>	<p>Please submit: A COMPLETED PRIOR AUTHORIZATION FORM (see link below)</p> <p>SYNAGIS PRIOR AUTHORIZATION AND PRESCRIPTION FORM</p> <p>To view the most up to date AAP Synagis Guidelines, follow the link below: AAP SYNAGIS GUIDELINES</p>	SYNAGIS PI
Synribo (omacetaxine)	<p>Indicated to treat adults with chronic phase (CP) or accelerated phase (AP) CML with resistance and/or intolerance to two or more TKIs.</p>	<p>Rx by Oncologist</p>	SYNRIBO PI
Syprine (trientine hydrochloride)	<p>Indicated in the treatment of patients with Wilson’s disease who are intolerant of penicillamine.</p>		SYPRINE PI
Tabrecta (capmatinib)	<p>Indicated for treatment of adults with metastatic NSCLC whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an approved test.</p>	<p>Rx by Oncologist</p>	TABRECTA PI
Tafinlar (dabrafenib)	<p>Indicated for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E mutation as detected by an FDA-approved test. TAFINLAR in combination with trametinib is indicated for the treatment of patients with unresectable or metastatic melanoma with BRAF V600E or V600K mutations as detected by an FDA-approved test.</p>	<p>Rx by Oncologist</p>	TAFINLAR PI
Tagrisso (osimertinib)	<p>Indicated for:</p> <ol style="list-style-type: none"> the treatment of first-line treatment of patients with metastatic NSCLC whose tumors have epidermal growth factor receptor (EGFR) Exon 19 deletions or exon 21 L858R mutations, as detected by an FDA approved test. the treatment of patients with metastatic EGFR T790M mutation- positive NSCLC, as detected by an FDA approved test, whose disease has progressed on or after EGFR TKI therapy. 	<p>Rx by Oncologist</p>	TAGRISSO PI
Talzenna (talazoparib)	<p>Indicated for the treatment of adult patients with deleterious or suspected deleterious germline BRCA-mutated (gBRCAm) HER2-negative locally advanced or metastatic breast cancer.</p>	<p>Rx by Oncologist</p>	TALZENNA PI
Tarceva (erlotinib)	<p>Indicated for:</p> <ol style="list-style-type: none"> treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA- approved test receiving 	<p>Rx by Oncologist</p>	TARCEVA PI

	<p>first-line, maintenance, or second or greater line treatment after progression following at least one prior chemotherapy regimen.</p> <p>2. first-line treatment of patients with locally advanced, unresectable or metastatic pancreatic cancer, in combination with gemcitabine.</p>		
Tasigna (nilotinib)	<p>Indicated for:</p> <ol style="list-style-type: none"> adult and pediatric patients greater than or equal to 1 year of age with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase. adult patients with chronic phase (CP) and accelerated phase (AP) Ph+ CML resistant to or intolerant to prior therapy that included imatinib. pediatric patients greater than or equal to 1 year of age with Ph+ CML-CP resistant or intolerant to prior tyrosine-kinase inhibitor (TKI) therapy. 	Rx by Oncologist	TASIGNA PI
Tavalisse (fostamatinib disodium hexahydrate)	Indicated for the treatment of thrombocytopenia in adult patients with chronic immune thrombocytopenia (ITP) who have had an insufficient response to a previous treatment.	Rx by Hematologist	TAVALISSE PI
Taytulla (norethindrone/ ethinyl estradiol capsules and ferrous fumarate)	See Oral Contraceptive	ANY OCP on prior authorization requires documentation demonstrating a compelling reason why formulary OCPs cannot be used [ex: intolerance, prior side effects, failures, etc. documented after a 3-month trial of formulary OCPs]	TAYTULLA PI
Tazverik (tazemetostat)	<p>Indicated for:</p> <ol style="list-style-type: none"> Adults and pediatric patients aged 16 years and older with metastatic or locally advanced epithelioid sarcoma not eligible for complete resection. Adult patients with relapsed or refractory follicular lymphoma whose tumors are positive for an EZH2 mutation as detected by an FDA- approved test and who have received at least 2 prior systemic therapies. Adult patients with relapsed or refractory follicular lymphoma who have no satisfactory alternative treatment options. 	Rx by Oncologist	TAZVERIK PI
Tibsovo (ivosidenib tablets)	<p>Indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with a susceptible IDH1 mutation as detected by an FDA-approved test in:</p> <ol style="list-style-type: none"> Adult patients with newly diagnosed AML who are > 75 years old or who have comorbidities that preclude use of intensive induction chemotherapy. Adult patients with relapsed or refractory AML. 	Rx by Oncologist	TIBSOVO PI
Trikafta (elexacaftor, ivacaftor, and tezacaftor)	Indicated for the treatment of cystic fibrosis (CF) in patients aged 12 years and older who have at least one F508del mutation in the CFTR gene. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to confirm the presence of at least one F508del mutation.	Rx by Pulmonologist	TRIKAFTA PI
Trodelyv (sacituzumab govitecan- hziy)	Indicated for the treatment of adult patients with metastatic triple-negative breast cancer who have received at least two prior therapies for metastatic disease.	Rx by Oncologist	TRODELVY PI

Truseltiq (infigratinib)	Indicated for the treatment of previously treated, unresectable locally advanced or metastatic cholangiocarcinoma (CCA) with a fibroblast growth factor receptor II (FGFR2) fusion in adult patients.	Rx by Oncologist	TRUSELTIQ PI
Tukysa (tucatinib)	Indicated in combination with trastuzumab and capecitabine for treatment of adult patients with advanced unresectable or metastatic HER2-positive breast cancer, including patients with brain metastases, who have received one or more prior anti-HER2-based regimens in the metastatic setting.	Rx by Oncologist	TUKYSA PI
Turalio (pepidartinib)	Indicated for the treatment of adult patients with symptomatic tenosynovial giant cell tumor associated with severe morbidity or functional limitations and not amenable to improvement with surgery.		TURALIO PI
Tykerb (lapatinib)	Indicated in combination with capecitabine for the treatment of patients with advanced or metastatic breast cancer whose tumors overexpress human epidermal growth factor receptor 2 (HER2) and who have received prior therapy including an anthracycline, a taxane, and trastuzumab. Patients should have disease progression on trastuzumab prior to initiation of treatment with TYKERB in combination with capecitabine.	Rx by Oncologist	TYKERB PI
Ubrelvy (ubrogepant)	Indicated for the acute treatment of migraine with or without aura in adults.	Member must have tried and failed NSAIDs and Triptans or have a contraindication to taking either of these medications.	UBRELVY PI
Venclexta (venetoclax)	Indicated: 1. The treatment of adult patients with chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL). 2. In combination with azacitidine, or decitabine, or low dose cytarabine for the treatment of newly diagnosed acute myeloid leukemia (AML) in adults 75 years or older, or who have comorbidities that preclude use of intensive induction chemotherapy.	Rx by Oncologist	VENCLEXTA PI
V-Go	Wearable insulin device indicated for use in adult patients requiring insulin.	Rx by Endocrinologist. Please click link below for External Insulin Pump Policy: MFC External Insulin Pumps Policy	V-GO WEBSITE
Viltepsa (viltolarsen)	Indicated for the treatment of Duchenne muscular dystrophy (DMD) in patients who have a confirmed mutation of the DMD gene that is amenable to exon 53 skipping. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	VILTEPSO PI
Vittrakvi (larotrectinib)	Indicated for the treatment of adult and pediatric patients with solid tumors that: 1. have a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known	Rx by Oncologist	VITRAKVI PI

	<p>acquired resistance mutation, and</p> <ol style="list-style-type: none"> are metastatic or where surgical resection is likely to result in severe morbidity, and have no satisfactory alternative treatments or that have progressed following treatment. 		
Vizimpro (dacomitinib)	Indicated for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test.	Rx by Oncologist	VIZIMPRO PI
Vimizim (elosulfase alfa)	Indicated for patients with Mucopolysaccharidosis type IVA (MPS IVA; Morquio A syndrome). Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.	***REQUIRES MFC PHYSICIAN REVIEW	VIMIZIM PI
Vocabria (cabotegravir)	Vocabria (cabotegravir) - Indicated in combination with EDURANT (rilpivirine) for short-term treatment of HIV-1 infection in adults who are virologically suppressed (HIV-1 RNA less than 50 copies/mL) on a stable antiretroviral regimen with no history of treatment failure and with no known or suspected resistance to either cabotegravir or rilpivirine, for use as: <ol style="list-style-type: none"> oral lead-in to assess the tolerability of cabotegravir prior to administration of CABENUVA (cabotegravir; rilpivirine) extended-release injectable suspensions. oral therapy for patients who will miss planned injection dosing with CABENUVA. 		VOCABRIA PI
Xadago (safinamide)	Indicated as adjunctive treatment to levodopa/carbidopa in patients with Parkinson's disease (PD) experiencing "off" episodes.	Rx by Neurologist	XADAGO PI
Xalkori (crizotinib)	Indicated for the treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) or ROS1-positive as detected by an FDA-approved test.	Rx by Oncologist	XALKORI PI
Xgeva (denosumab)	Indicated for: <ol style="list-style-type: none"> Prevention of skeletal-related events in patients with multiple myeloma and in patients with bone metastases from solid tumors. Treatment of adults and skeletally mature adolescents with giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity. Treatment of hypercalcemia of malignancy refractory to bisphosphonate therapy. 	Rx by Oncologist	XGEVA PI
Xiidra (lifitegrast ophthal)	Indicated for the treatment of the signs and symptoms of dry eye disease.	Must have tried and failed artificial tears.	XIIDRA PI
Xolair (omalizumab)	Indicated for: <ol style="list-style-type: none"> moderate to severe persistent asthma in patients 6 years of age and older with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms that are inadequately controlled with inhaled corticosteroids. chronic idiopathic urticaria in adults and adolescents (12 years of age and above) who remain symptomatic despite H1 antihistamine treatment. 	Rx by Allergist or Pulmonologist Regarding ASTHMA indication only: <ol style="list-style-type: none"> moderate to severe persistent ALLERGIC asthma (confirmed by a positive skin test or RAST for ≥ 1 perennial aeroallergen). 	XOLAIR PI

		<ol style="list-style-type: none"> 2. IgE level obtained prior to initiation of therapy. 3. currently using an inhaled corticosteroid at maximum dose; compliance must be confirmed in the patient's Caremark profile. 4. currently using a long-acting inhaled beta2-agonist OR a leukotriene modifier; compliance must be confirmed in the patient's Caremark profile. 	
Xospata (gilteritinib)	Indicated for the treatment of adult patients who have relapsed or refractory acute myeloid leukemia with a FLT3 mutation as detected by an FDA-approved test.	Rx by Oncologist	XOSPATA PI
Xpovio (selinexor)	Indicated for in combination with dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least four prior therapies and whose disease is refractory to at least two proteasome inhibitors, at least two immunomodulatory agents, and an anti-CD38 monoclonal antibody.	Rx by Oncologist	XPOVIO
Xyrem (sodium oxybate)	<p>Indicated for the treatment of cataplexy or excessive daytime sleepiness (EDS) in patients 7 years of age and older with narcolepsy.</p> <p>***Xyrem may only be dispensed to patients enrolled in the Xyrem Success Program</p>	<ol style="list-style-type: none"> 1. patient > 16 years old 2. alternative diagnoses must have been excluded 3. for cataplexy, must have failed tricyclic or SSRIs 4. for excessive daytime sleepiness, must have failed at least one formulary stimulant treatment (ex: methylphenidate or dextroamphetamine) 5. initial approval for maximum of 1-month supply with subsequent renewals for maximum approval period of 3 months at a time (Patients are to be re-evaluated by physician no less frequently than every 3 months) 6. Rx by Neurologist 	XYREM PI
Yescarta (axicabtagene ciloleucel)	<p>Indicated for the treatment of adult patients with relapsed or refractory large B- cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, primary mediastinal large B-cell lymphoma, high grade B-cell lymphoma, and DLBCL arising from follicular lymphoma.</p> <p>In accordance with criteria developed by the Maryland Medicaid Program, MedStar Family</p>	Rx by Oncologist	YESCARTA PI

	<p>Choice considers Yescarta (Axicabtagene Ciloleucel) medically necessary when ALL of the following criteria are met:</p> <ol style="list-style-type: none"> 1. Recipient is 18 years of age or older; AND 2. Histologically confirmed diagnosis of one of the following types of aggressive non-Hodgkin’s lymphoma <ol style="list-style-type: none"> a. Diffuse large B-cell lymphoma (DLBCL), not otherwise specified; or b. High-grade B-cell lymphoma; or c. Primary mediastinal large B-cell lymphoma; or d. Transformed follicular lymphoma; AND 3. Relapsed or refractory disease, when <ol style="list-style-type: none"> a. Recipient has previously received two or more lines of systemic therapy; and b. Disease is refractory to the most recent therapy or relapsed within 1 year after autologous hematopoietic stem cell transplantation (HSCT); AND 4. Must have received adequate prior therapy including, at a minimum, all of the following: <ol style="list-style-type: none"> a. An anthracycline-containing chemotherapy regimen; and b. For CD20+ disease, anti-CD20 monoclonal antibody; and c. For subjects with transformed follicular lymphoma, prior chemotherapy for follicular lymphoma with chemotherapy refractory disease after transformation to DLBCL; AND 5. Documentation of all of the following clinical findings: <ol style="list-style-type: none"> a. Eastern Cooperative Oncology Group (ECOG) performance status of 0-1; and b. Adequate cardiac, pulmonary, and other organ function (as determined by protocol from treatment facility); AND 6. The treatment facility that dispenses and administers Yescarta is enrolled and complies with the Risk Evaluation and Mitigation Strategy; AND 7. One-time, single administration with dosing in accordance with the FDA label <p>Yescarta (Axicabtagene ciloleucel) is considered investigational and not medically necessary when the above medically necessary criteria are not met, and for all other indications, including but not limited to:</p> <ol style="list-style-type: none"> 1. History of malignancy other than nonmelanoma skin cancer or carcinoma in situ (e.g. cervix, bladder, breast) or follicular lymphoma unless disease free for at least 3 years; or 2. Any central nervous system (CNS) disease, for example, detectable CSF malignant cells, brain metastases, CNS lymphoma, or a history or presence of CNS disorders such as seizure disorder, cerebrovascular ischemia/hemorrhage, dementia, cerebellar disease, or autoimmune disease with CNS involvement ; or 3. History of allogeneic stem cell transplant, chimeric antigen receptor therapy or other genetically modified T-cell therapy; or 4. Active, uncontrolled infection; or Human immunodeficiency virus (HIV); or Hepatitis B or C (if viral load is detectable). 		
<p>Zejula (niraparib)</p>	<p>Indicated for:</p> <ol style="list-style-type: none"> 1. the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial 	<p>Rx by Oncologist</p>	<p>ZEJULA PI</p>

	<p>response to platinum-based chemotherapy.</p> <ol style="list-style-type: none"> the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to first-line platinum-based chemotherapy. the treatment of adult patients with advanced ovarian, fallopian tube, or primary peritoneal cancer who have been treated with three more prior chemotherapy regimens and whose cancer is associated with homologous recombination deficiency (HRD) positive status defined by either: <ul style="list-style-type: none"> a deleterious or suspected deleterious BRCA mutation, or genomic instability and who have progressed more than six months after response to the last platinum-based chemotherapy. 		
Zelboraf (vemurafenib)	<p>Indicated for:</p> <ol style="list-style-type: none"> the treatment of patients with unresectable or metastatic melanoma with BRAF V600E mutation as detected by an FDA-approved test. the treatment of patients with Erdheim Chester Disease with BRAF V600 mutation. 	Rx by Oncologist or Dermatologist	ZELBORAF PI
Zoladex (goserelin)	<p>Indicated for:</p> <ol style="list-style-type: none"> palliative treatment of advanced carcinoma of the prostate. (3.6 and 10.8mg) use in combination with flutamide for the management of locally confined stage T2b-T4 (Stage B2-C) carcinoma of the prostate. (3.6 and 10.8 mg) management of endometriosis. (3.6 mg) palliative treatment of advanced breast cancer in pre- and peri-menopausal women. (3.6 mg) use as an agent to cause endometrial thinning agent prior to endometrial ablation for dysfunctional uterine bleeding. (3.6 mg) for the management of endometriosis, including pain relief and reduction of endometriotic lesions for the duration of therapy. Experience with ZOLADEX for the management of endometriosis has been limited to women 18 years of age and older treated for 6 months. 	Rx by Oncologist	ZOLADEX 3.6 mg PI ZOLADEX 10.8 mg PI
Zolgensma (onasemnogene abeparvovec-xioi)	<p>Indicated for the treatment of pediatric patients less than 2 years of age with spinal muscular atrophy (SMA) with bi-allelic mutations in the survival motor neuron 1 gene.</p> <p>Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.</p>	***REQUIRES MFC PHYSICIAN REVIEW	ZOLGENSMA PI
Zontivity (vorapaxar)	Indicated for the reduction of thrombotic cardiovascular events in patients with a history of myocardial infarction or with peripheral arterial disease.	Rx by Cardiology, Neurology or Vascular Surgery	ZONTIVITY PI
Zurampic (lesinurad)	Indicated in combination with a xanthine oxidase inhibitor for the treatment of hyperuricemia associated with gout in patients who have not achieved target serum uric acid levels with a xanthine oxidase inhibitor alone.	Rx by Rheumatologist	ZURAMPIC PI

Zydelig (idelalisib)	Indicated for: <ol style="list-style-type: none"> 1. treatment of patients with relapsed chronic lymphocytic leukemia, in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities. 2. treatment of patients with relapsed follicular B-cell non-Hodgkin lymphoma in patients who have received at least two prior systemic therapies. 3. treatment of patients with relapsed small lymphocytic lymphoma in patients who have received at least two prior systemic therapies. 	Rx by Oncologist	ZYDELIG PI
Zykadia (ceritinib)	Indicated for the treatment of adults with metastatic non-small cell lung cancer whose tumors are anaplastic lymphoma kinase-positive as detected by an FDA-approved test.	Rx by Oncologist	ZYKADIA PI
Zynlonta (loncastuximab tesirine-lpyl)	Indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, DLBCL arising from low-grade lymphoma, and high-grade B-cell lymphoma.	Rx by Oncologist	ZYNLONTA PI